

United States Patent [19]

Kortright et al.

[11] Patent Number: 4,752,563

[45] Date of Patent: Jun. 21, 1988

- [54] MONOCLONAL ANTIBODY FOR RECOVERY OF LEUKOCYTES IN HUMAN PERIPHERAL BLOOD AND METHOD OF RECOVERY EMPLOYING SAID MONOCLONAL ANTIBODY
- [75] Inventors: Kenneth H. Kortright, Copper City; David E. Hofheinz, Homestead, both of Fla.
- [73] Assignee: Coulter Corporation, Hialeah, Fla.
- [21] Appl. No.: 799,489
- [22] Filed: Nov. 19, 1985
- [51] Int. Cl.⁴ C12Q 1/02; G01N 33/577
- [52] U.S. Cl. 435/2; 435/4; 435/7; 435/29; 435/68; 435/70; 435/172.2; 435/240.27; 436/518; 436/526; 436/528; 436/534; 436/548; 530/387; 935/108
- [58] Field of Search 435/2, 7, 29, 4, 68, 435/70, 172.2, 240, 241; 436/548, 518, 526, 528, 534; 935/103, 104, 108, 110; 530/387

[56] References Cited

U.S. PATENT DOCUMENTS

- 4,429,008 1/1984 Martin 428/402.2
4,598,051 7/1986 Papahadjopoulos 436/512

OTHER PUBLICATIONS

- Robinson, J. et al., Nature, 289:68-71, (1-1981).
Anstee, D. J. et al., Eur. J. Immunology, 12:228-232, (1982).
Barsaum, A. H. et al., Molecular Immunology, 22(4):361-367, (1985).
Rearden, A. et al., Molecular Immunology, 22(4):369-378, (1985).

Langlois, R. G. et al., Journal of Immunology, 134(6):4009-4017, (6-1985).
Bigbee, W. L. et al., Journal of Immunology, 133(6):3149-3155, (12-1984).
Bigbee, W. L. et al., Molecular Immunology, 20(12):1351-1362, (1983).
Liszka, K. et al., Amer. J. Hematology, 15:219-226, (1983).
Ridgewell, K. et al., Biochem. Journal, 209:273-276, (1983).
Ochiai, Y. et al., J. of Immunology, 131:864-868, (8-1983).
Bragman, K. S. et al., Biochem. Biophys. Acta, 730:187-195, (1983).

Primary Examiner—Margaret Moskowitz
Attorney, Agent, or Firm—Myron C. Cass

[57] ABSTRACT

A murine monoclonal antibody which selectively binds to the determinant site or specific epitope of glycoprotein A exposed on the membrane of the erythrocyte and does not bind to other glycoproteins. The monoclonal antibody is coated on a microsphere or substrate of a suitable monodispersed variety and utilized in a separation procedure for recovery of white blood cell subsets without lysing of erythrocytes. The bound microspheres are recovered without adverse depletion of the white blood cell population of the sample. The microsphere or bead-to-cell ratio employed is most acceptable for commercial applications of the invention. The invention thus enables accurate assaying of white blood cell subsets in circulating peripheral blood without resorting to lysing of erythrocytes.

8 Claims, No Drawings